



CHINA 中国地质(英文)
GEOLOGY



China Geological Survey conducted the first natural gas hydrates production test in the South China Sea

Geometry and kinematics of brittle deformation in the Central Cameroon Shear Zone (K é kem area): Implication for gold exploration within the Central Africa Fold Belt in Cameroon

De Pesquidoux I Tchaptchet Tchato, Jacqueline Tchakounte, Aur é lie Ngamy Kamwa, Jean Pierre Tchouankoue, Soumyajit Mukherjee

Citation: De Pesquidoux I Tchaptchet Tchato, Jacqueline Tchakounte, Aur é lie Ngamy Kamwa, Jean Pierre Tchouankoue, Soumyajit Mukherjee, 2021. Geometry and kinematics of brittle deformation in the Central Cameroon Shear Zone (K é kem area): Implication for gold exploration within the Central Africa Fold Belt in Cameroon, *China Geology*, 4, 245–255. doi: [10.31035/cg2020058](https://doi.org/10.31035/cg2020058).

View online: <https://doi.org/10.31035/cg2020058>

Related articles that may interest you

Deep gold mineralization features of Jiaojia metallogenic belt, Jiaodong gold Province: Based on the breakthrough of 3000 m exploration drilling

China Geology. 2020, 3(3), 385 <https://doi.org/10.31035/cg2020048>

LA–ICP–MS *in situ* analyses of the pyrites in Dongyang gold deposit, Southeast China: Implications to the gold mineralization

China Geology. 2020, 3(2), 230 <https://doi.org/10.31035/cg2018123>

An integrated ore prospecting model for the Nyasirori gold deposit in Tanzania

China Geology. 2019, 2(4), 407 <https://doi.org/10.31035/cg2018127>

Effects of heavy metal pollution on farmland soils and crops: A case study of the Xiaoqingling Gold Belt, China

China Geology. 2020, 3(3), 402 <https://doi.org/10.31035/cg2020024>

China has launched a deep gold prospecting demonstration project to evaluate gold resource potential within 3000 m underground in the east of the North China Craton

China Geology. 2018, 1(4), 572 <https://doi.org/10.31035/cg2018047>

Types of uranium deposits in central Zhuguang Mountains in Hunan Province, South China and their metallogenic regularity and prospecting directions

China Geology. 2020, 3(3), 411 <https://doi.org/10.31035/cg2020040>



Geometry and kinematics of brittle deformation in the Central Cameroon Shear Zone (Kékem area): Implication for gold exploration within the Central Africa Fold Belt in Cameroon

De Pesquidoux I Tchaptchet Tchato^a, Jacqueline Tchakounte^a, Aurélie Ngamy Kamwa^b, Jean Pierre Tchouankoue^a, Soumyajit Mukherjee^{c,*}

^a Department of Earth Sciences, Faculty of Sciences, University of Yaoundé I, Yaoundé P.O.BOX : 812, Cameroon

^b Department of Mining Engineering and Minerals processing, Faculty of Mines and Petroleum Industries, University of Maroua, Maroua P.O.BOX : 46, Cameroon

^c Geodynamics Laboratory, Department of Earth Sciences, Indian Institute of Technology Bombay, Powai, Mumbai Maharashtra 410076, India

ARTICLE INFO

Article history:

Received 15 April 2020

Received in revised form 28 June 2020

Accepted 29 July 2020

Available online 28 October 2020

Keywords:

Gold deposit
 Gold mineralization
 Gold exploration
 SRTM images
 Riedel fractures
 Simple shear
 Congo Craton
 Kékem
 Central Africa fold belt
 Cameroon

ABSTRACT

The Central Africa Fold Belt (CAFB) is a collision belt endowed with gold deposits in Eastern Cameroon area mined for about 50 years. However, favorable areas for gold exploration are poorly known. This paper presents (1) the kinematics of the brittle deformation in the Kékem area in the SW portion of the Central Cameroon Shear Zone and (2) constraints gold mineralization events with respect to the collisional evolution of the CAFB. The authors interpret that the conjugate ENE to E and NNW to NW trending lineament corresponds to the synthetic (R) and the antithetic (R') shears, which accompanied the dextral slip along the NE to ENE striking shear. The latter coincides with the last 570–552 Ma D₃ dextral simple shear-dominated transpression, which is parallel to the Bétaré Oya shear zone hosting gold deposits. Gold mineralizations, which mainly occurred during the last dextral shearing, are disseminated within quartz veins associated to Riedel's previous structures reactivated due to late collisional activities of the CAFB as brittle deformation. Gold mineralizations occurred mainly during the 570–552 Ma D₃ event. The reactivation, which might be due to dextral simple shear during mylonitization, plausibly remobilized the early gold deposits hosted in syn-compressional rocks and/or possibly focused deep-sourced fluid mixed with those released by dehydration. Therefore, the Central Cameroon Shear Zone where Kékem is located, and which shows similar petrographical and structural features to those controlling Batouri gold district, is a target area for gold exploration in Cameroon.

©2021 China Geology Editorial Office.

1. Introduction

The Central African Fold Belt (CAFB) in Cameroon (Toteu SF et al., 2001), a part of the vast Brasiliano-Pan-African Fold Belt (Fig. 1a) in the pre-drift reconstruction (Archanjo CJ et al., 2013), is a typical collisional belt developed in a subduction-collisional setting (Owona S et al., 2010; Bouyo MH et al., 2013). Its evolution is characterized by an early crustal thickening followed by transpression with a sinistral- and then by a dextral shear along about the NE to ENE striking Central Cameroon Shear Zone (CCSZ; Ngako V

et al., 1991) and the Bétaré Oya Shear zone (Kankeu B et al., 2009). A complex network of fault system characterized by a N70°E parallel en-echelon shear zones, inter-connected by a N40°E directed S-type restraining bend exists in the Magba area (Njonfang E et al., 2008). This S-type bend, extending southward up to the Foutoni and the Kékem areas (Tchaptchet TD et al., 2009), is associated with a network of secondary fault and fractures around Fouban (Njonfang E et al., 2008), Kékem (Tchaptchet TD, 2017) and the Tombel plain (Njome MS and Suh CE, 2005). Although Simeni NAW et al. (2017) linked secondary fractures with the Pan-African structures, their geometry and kinematics remained poorly documented. Interestingly, some of these transpressional shear zones (Tcholliré Shear Zone: TSZ; Bétaré Oya Shear Zone: BOSZ) and associated subsidiary shears in their adjacent areas host gold deposits (Suh CE et al., 2006; Tchameni V et al., 2013).

First author: E-mail address: depesquidoux@yahoo.fr (De Pesquidoux I Tchaptchet Tchato).

* Corresponding author: E-mail address: soumyajitm@gmail.com (Soumyajit Mukherjee).

doi:10.31035/cg2020058

2096-5192/© 2021 China Geology Editorial Office.

E.g. vein-type gold deposits (0.04 g/t to 30 g/t and even 66 g/t Au) are associated to the NE-SW trending shear zone within the 620 Ma metagranite and in altered wall rock up to 18.5 g/t Au (Asaah VA et al., 2014; Suh CE et al., 2006, Vishiti A et al., 2019) are reported in Batouri and adjacent areas, which are affected by the Betaré Oya Shear Zone (Kankeu B et al., 2009). The reserve of this gold which is mined in Cameroon for more than 50 years (Asaah VA et al., 2014) is estimated to be at least 15 t (Mélisi JP et al., 2006).

This study performs geometric and kinematic analyses of the lineaments in the Kékem area, located at SW of the CCSZ. These analyses are based on (1) lineaments identification by combining geomorphological parameter, (2) digital elevation model, and (3) field data. CCSZ parallels the gold-enriched BOSZ. The gold mineralization processes and their plausible timing with respect to the tectonic evolution of the CAFB are discussed. The Kékem area, located SW of the CCSZ is assessed for the suitability for gold exploration following its comparison with the Betaré Oya Shear Zone. The area

displays similar tectonic evolution and lithological features that apparently controlled gold mineralization at the BOSZ.

2. Regional geology

The Kékem area (Fig. 1b) is located at the SW of the about NE striking CCSZ (Ngako V et al., 1991; Njonfang E et al., 2008; Tchaptchet TD et al., 2009) of the CAFB in Cameroon (Fig. 1b). The CAFB is divided into three geodynamic domains (Toteu SF et al., 2004). (1) The northern domain is made up of the Poli, the Léré and the Mayo Kebbi meta-sedimentary series all intruded by pre-, syn- and post-tectonic (750–680 Ma) calc-alkaline rocks emplaced in an Early Neoproterozoic back-arc or a continental magmatic arc basin (Toteu SF et al., 2006; Bouyo MH et al., 2015). These intrusions define an NNE-SSW corridor marking the southern limit of the northern domain. (2) The southern domain is monocyclic with granulitic nappes made up of Neoproterozoic metasedimentary units (Mbalmayo, Yaounde and Ntui-Betamba). These units, consisting of schists, garnet-kyanite-




Fig. 1. a–Pan-African shear zone network in pre-Mesozoic reconstruction (after Cabry R, 1989). SZ = Shear Zone: C = Cameroon area. b–Pan-African structural map of Cameroon from Njonfang E et al. (2008) and location of the study area. 1–Quaternary sediments; 2–Cameroon Line volcanism; 3–Cameroon Line plutonism; 4–Mesozoic sediments (Benue through); 5–Late syntectonic sub-alkaline granitoids; 6–Lom syntectonic basin (meta-sediments, conglomerates, volcanic ashes and lavas); 7–Western Cameroon Domain (WCD: Early syntectonic basic to intermediate calc-alkaline intrusions, 660–600 Ma); 8a–Poli Group (active margin Neoproterozoic supracrustal and juvenile intrusions); 8b–Yaounde Group (intracratonic deposits); 9–Massenya-Ounianga gravity highs (10–30 mGal); 10–Adamawa-Yadé and Nyong Paleoproterozoic remnants; 11–craton and inferred craton; 12– S_2 foliation and L_2 lineation trends; 13– F_2 upright and overturned antiforms; 14–syn- D_2 main frontal thrust zone; 15–syn- D_1 thrust zone (separates the LP to MPzone in the North from the HP zone in South); 16–syn- D_3 sense of shear movement; 17–syn- D_2 sense of shear movement. Large grey arrow:syn- $D_{1,3}$ regional main stress direction. Thick lines-shear zones (SZ): BSZ–Balché SZ; BNMB = Buffle Noir - Mayo Baléo SZ; CCSZ–Central Cameroon SZ; GGSZ– Godé - Gormaya SZ; MNSZ–Mayo Nolti SZ; RLSZ–Rocher du Loup SZ; SSZ–Sanaga SZ; Ma–Magha; Wa–Wakaa. Small squares: BA–Bandja complex.

bearing mica-schists and garnet-kyanite gneisses, formed originally in either a passive margin (Mvondo H et al., 2003) or an active margin (Standal H et al., 2006) at the northern edge of the Congo Craton. They are associated with alkaline magmatic rocks and the entire unit thrusts onto the Archean Congo Craton towards south (Ngako V et al., 2008).

The Pan-African tectonics is characterized by three main deformation phases. The significance of these deformations is interpreted differently. According to Mvondo H et al. (2007), the southern domain is characterized by two compressive events, D_1 and D_3 , with E-W to NW-SE maximum stress direction. These two phases are separated by an intermediate D_2 event, of extension parallel to the orogen synchronous with the peak metamorphism. This evolution is consistent with a magmatic-arc collision setting (Mvondo H et al., 2007). D_1 and D_3 are compressive tectonics with nappe stacking during D_1 - D_2 interphase coeval with the 620 ± 10 Ma granulitic metamorphic peak and anatexis (Penaye J et al., 1993) defining a suture zone following collision between the Congo Craton to the south and the Paleoproterozoic crust to the north. These two domains are separated by Adamawa-Yadé domain, which is transected by the CCSZ and is sandwiched between the $N70^\circ E$ striking Tcholliré-Banyo Shear Zone (TBSZ) to the N and the Sanaga Shear Zone (SSZ) to the south (Fig. 1b; Toteu SF et al., 2004). These transcurrent CCSZ faults are regarded as possible continuation of the major shear zones of NE Brazil in the pre-drift Gondwana reconstruction (Archanjo CJ et al., 2013). This polycyclic domain is made up of an assemblage of about 2100 Ma remnants of Paleoproterozoic rock with an Archean heritage (Penaye J et al., 2004; Toteu SF et al., 2001; Ganwa AA et al., 2016) and Neoproterozoic meta-sedimentary rocks recrystallized in upper amphibolite to granulites facies metamorphism during 630–600 Ma of crustal thickening (Tchaptchet TD et al., 2009; Tchaptchet TD, 2011; Bouyo MH et al., 2013). Recent works evidence that the Adamawa domain corresponds an orthoderived Archean (2.9–2.5 Ga) cratonic nucleus having TTG composition (Tchakounté et al., 2017). These formations are further intruded by Pan-African granitoids whose emplacement age in Kékem is about 576–560 Ma (Kwékam M et al., 2013, Tchaptchet TD, 2011) and 640–620 Ma at Batouri gold district (Asaah VA et al., 2014). Subsequently in this article D_4 phase will denote the brittle deformation.

The Pan-African tectonic evolution of this domain is characterized by an early D_1 deformation (630–600 Ma; Bouyo MH et al., 2013) of crustal thickening followed by 620–585 Ma D_2 phase (Ngako V and Njonfang E, 2011) with N-S to ENE-WSW striking sinistral transcurrent/transpressional shear zones (e.g., the Mayo Nolti shear zone, the Gordé Gormaya shear zone and the Rocher du Loup shear zone: Ngako V et al., 2008) developed during amphibolite facies metamorphism (Njonfang E et al., 2008). The about 585–583 Ma D_3 event (Ngako V et al., 2008) is marked by a about NE to ENE trending dextral strike slip faults (e.g., CCSZ; Fig. 1b; Ngako V et al., 2008) that transect the central domain- locally

known as the Fouban Shear Zone (FSZ; Njonfang E et al., 2008) or the Fotouni-Fouban Shear Zone (FFSZ; Tcheumenak Kouémo J et al., 2014). It is about 500 km long and 5–12 km wide (Njonfang E et al., 2008).

The Kékem area (Fig. 2), at the SW of the central domain, comprises Paleoproterozoic magma-derived migmatitic gneiss (Dumort JC, 1968; Tchaptchet TD 2009), displaying 2.7–1.5 Ga (Sm/Nd TDM age) Archean to Paleoproterozoic ages (Penaye J et al., 1993; Toteu SF et al., 2001; Tchaptchet TD, 2011). The rocks consist of garnet-bearing pyroxene gneiss dated at 2100 Ma (U/Pb on zircon; Penaye J et al., 1993; Toteu SF et al., 2001) and biotite-amphibole gneiss. The rocks recrystallized at granulites facies metamorphic conditions and subsequently retrogressed into amphibolite facies during the Pan-African orogeny (Penaye J et al., 2004). They are associated with lenses of Neoproterozoic metasedimentary rocks, garnet-biotite gneiss and sillimanite-garnet-biotite gneiss (Tchaptchet TD et al., 2009) that recrystallized under the upper amphibolites-facies metamorphic condition about 580 Ma (U-Th-Pb on monazite; Tchaptchet TD et al., 2009) or 600 Ma (U-Pb on zircon; Tchaptchet TD, 2011) during crustal thickening, following a continental collision (Tchaptchet TD et al., 2009). These migmatitic gneisses have been intruded by high-K calc-alkaline to shoshonitic post-collisional magmatic rocks (biotite granite with enclave of diorite, amphibole granite, muscovite-biotite granite, gabbro and norites) at 570–560 Ma (Dumort JC, 1968; Tchaptchet TD et al., 2009, 2011; Kwékam M et al., 2013). This magmatism is coeval with the activity of the $N40^\circ E$ striking high temperature shear zone constrained at 570–552 Ma (U-Th-Pb dating of monazite (Tchaptchet TD et al., 2009).

The shear zone is associated to secondary fractures filled in by sheet-like granite and tholeiitic basalt dykes originated from the melting of subcontinental lithospheric mantle by early fragmentation of the West Gondwana (Tchaptchet TD et al., 2017). Migmatitic gneisses and magmatic rocks are covered partially by the 10–5 Ma alkaline basalts (Tchuimegnie Ngongang NB et al., 2015) of the Cameroon Volcanic Line.

3. Methodology

This work uses combined methodologies of Ganwa AA et al. (2007) and Akam JM et al. (2014) to identify lineaments. According to Ganwa AA et al. (2007), the analysis of geomorphologic parameters is very useful for identifying lineaments in the Pan-African setting. River flow can be tectonic controlled (Misra AA et al., 2014), and hence, can help in deciphering lineaments. Not all drainage patterns are tectonics-controlled. Any specific drainage pattern (e.g., the rectangular pattern) was not observed in the study area (Dasgupta S and Mukherjee, 2019). The hydrological network as in the 1 : 50000 scale topo-sheet (Fig. 3) was used and the lineaments identified are shown as black lines in Fig. 3. Then, this first lineament map is superposed on a second one obtained from digital elevation model of the Shuttle Radar




Fig. 2. Sketch of the geological map of the study area, extracted from Dumort JC (1968).




Fig. 3. Topographic map of the studied area with lineament identified from hydrographic network. This figure is generated from the SRTM DEM. Alternately, hillshade data derived from DEM may also be used.

Topography Mission (SRTM) following the methodology of Akam JM et al. (2014). This operation is based on 2001 (SRTM_ffB03_p186r057) satellite images (downloaded from internet and expressed in 3D on previously geo-referenced digital data; Fig 4). This SRTM DEM has a better resolution and image enhancement entails selection of subset of information to be displayed (Neawsuparp K and Charusir P, 2004). The shaded relief method, as an enhancement technique, discriminates structural features indicated by homogeneously colored area, corresponding to lineaments that are different from the background colour. These lineaments are extracted manually, and only the natural lineaments are taken. This operation is performed by using the Global Mapper software (version 10). Fig. 4 is superposed on Fig. 3 to obtain the lineament map of the study area (Fig. 5). In other words, lineaments extracted have been overlain on the SRTM DEM of the area. Only regional lineaments were selected for interpretation (Fig. 6a). SRTM images were downloaded and were worked on.

4. Results and discussion

4.1. Geometrical analyses

As per Fig. 6b, five fractures sets striking ENE, E, NW, NNE and NE are distinguished. The ENE fractures are most numerous (Fig.7a). Some of them deflect earlier foliation through dextral shear (Fig.7a). The E-W trending fractures are




Fig. 4. Digital elevation model from the Shuttle Radar Topography-Mission (SRTM) with identified lineaments. Black curved lines show curved foliations.




Fig. 5. Lineaments map from the hydrographic network and the SRTM image: Notice that red colored fractures are from hydrographic network and dark colored fractures from SRTM.

the second predominant that are commonly expose in the SE portion of the Kékém area (Fig. 7b). Some of them dextrally deflect the earlier foliation planes (Fig. 7c). A few sets of such fracture are filled with basalts (presumably dykes) and

granites in the Banwa locality (Figs. 7b, c). Few fractures cut across sheet like igneous bodies (Fig. 7d). The NNE-striking planes are less abundant and are marked by sinistral shear affecting the earlier foliations (Fig. 7e). The NE-trending fractures are least numerous (Fig. 7f). Fig. 8 presents the synoptic geometrical arrangement of these secondary fractures. The ENE fractures make about 15° with the NE and the E trending fracture sets. Moreover, ENE and NW trending fractures constitute conjugate sets at about 60°.

The geometry of the fault plane and the type of faulting depend on the internal angle of friction, strain rate, stress state (Ahlgren SG, 2001), vorticity and rheological properties of the rocks (Misra S et al., 2009; also see Mukherjee S and Khonsari, 2018). The complex fault pattern is also characterized by the Riedel conjugate set- comprising of synthetic Riedel fractures (R) and conjugate antithetic Riedel fractures (R'). This conjugate fracture set, most conspicuous as per the Anderson's model, is accompanied sometimes by synthetic P-shear fractures and tensional T fractures at about 45° with the primary shear Y-planes.

Fig. 8 is deduced based on the brittle planes' orientations for the Kékém area (Fig. 8). These plane are similar to those reported in the Bafia and Maham (Ngamy KA et al., 2016; Simeni NAW et al. 2017; Tchakounté J, 2020) SE of Kékém. The dextral shear along the NE to ENE striking shear zone is suggested to be the main fault. This implies that the NNE/NE, ENE, E-W and NW trending fractures correspond most plausibly to P, Y, R and R/in the complex fault pattern recorded in a strike slip shear zone (Simeni NAW et al., 2017).

4.2. Kinematics of fault-network in the Kékém area

As discussed in the previous section, the synoptic map of the Kékém area (Fig. 4) shows a fault pattern consistent with a right-handed Riedel shear with an ENE striking primary shear zone. This fault pattern matches with the dyke offsets of Simeni NAW et al. (2017) from the southern domain of the Cameroon Volcanic Line. The strike of the main shear zone coincides with that of the dextral regional CCSZ recorded geophysical from the basement rocks of the central domain (Noutchogwé TC et al., 2010).

Brittle deformation modeling shows that the maximum principal stress axis (σ_1) bisects the 60° angle between conjugate R and R' planes (Davis GH et al., 2000). Following this, σ_1 should trend E-W in this study area (Fig. 8). This orientation is compatible with the E-W to NW-SE maximum stress field direction ascribed to the D₃ folding in the southern domain (Mvondo H et al., 2003) or that of the NW-SE to WNW-ESE of the last dextral simple shear dominated transpression along the CCSZ in the central domain (Ngako V et al., 2008; Kankeu B et al., 2009; Ngako V and Njonfang E, 2011; Bella Nké BE et al., 2018) in the CAFB. The maximum stress field direction and the D₃ simple shear-dominated transpression are compatible, and this rotated the σ_1 -axis clockwise (Ngako V et al., 2008; Ngako V and Njonfang E,




Fig. 6. Synthetic lineaments map from the hydrographic network (a) and the SRTM image (b). Only regional lineaments selected.

2011; Bella Nké BE et al., 2018). The fractures system is coeval to the simple shear-dominated transpression. This hypothesis is consistent with Davis GH et al., (2000). As per Davis GH et al. (2000), shear zone experiencing simple shear is accompanied by Riedel's fractures kinematic model. Therefore, the reactivation of previous structures ascribed to D₃ transpression (Ngako V et al., 2008) can be envisaged for fracture systems recorded along the CCSZ (Njome MS and Suh CE, 2005) as well as the BOSZ (Takodjou WJD et al., 2016).

5. Implication for the evolution of the CCSZ in Cameroon

The CAFB (Cameroon) witnessed a tectonic indent collision between the East Sahara Block (ESB) and the Sao-Francisco - Congo and West African cratons (Ngako V et al., 2008). This collision is characterized by an early crustal thickening that existed during thrust and nappes tectonics (Ngako V et al., 2008; Owona S et al., 2010; Tchaptchet TD et al., 2009; Bouyo MH et al., 2013), followed by a 620–585 Ma sinistral shear (Ngako V et al., 2008; Tcheumenak Kouémo J et al., 2014) coeval with the emplacement of calc-alkaline to high K calc-alkaline granitoids. Finally, a D₃ dextral simple shear-dominated transpression (Ngako V et al., 2008, Njonfang E et al., 2008; Kankeu B et al., 2009; Bella Nké BE et al., 2018) at 585–538 Ma (Ngako V et al., 2008; Njonfang E et al., 2008) or 570–552 Ma (Tchaptchet TD et al., 2009) happened. These shear zones constitute a complex network of fault system characterized by a N70°E parallel encheelon pattern, presumably a “step-over” indicating intense shearing (reviewed in Mukherjee S 2013), inter-connected by a N40°E directed S-type restraining bend around Magba

(Njonfang E et al., 2008), which extend southward up to the Foutoni and the Kékem areas (Tcheumenak Kouémo J et al., 2014; Tchaptchet TD et al., 2009).

The reactivation characterizing the late collision (Ngako V et al., 2008) operate dominantly through simple shear (Bella Nké BE et al., 2018) inducing strike-slip (Misra S et al., 2009) and the Riedel model fractures. Simulation of faulting in zone of continental transpression by Schreurs G and Colletta B (1998) show that as deformation intensifies, the transverse component of the transpression increases the strike-slip leading to rocks failure. These fractures widespread along the CCSZ in Manjo (Njombe MS and Suh CE, 2005), Magba (Njonfang E et al., 2008), and the Betaré Oya shear zone (Som Mbang MC et al., 2018) are associated with sub-vertical sheet-like granitic veins and sub-continental mantle-derived tholeiitic basalt dykes characteristic of extension (Tchaptchet TD et al., 2017). Simple shear within the CCSZ could be the main mechanism that initiated the transition from ductile to brittle deformation marking the onset of extensional tectonics corresponding to the D₄ event. The later postdates the 570–552 Ma dextral shear along the CCSZ. This timing is compatible with the extensional tectonics-related 550 Ma alkaline granites within the fold belt (Toteu SF et al., 2001).

6. Tectonic setting and gold mineralization in the CAFB (Cameroon)

The CAFB in Cameroon is endowed with orogenic gold deposits (Tchameni R et al., 2013; Asaah VA et al., 2014) associated with compression and transpression. Indeed, the gold-bearing syenogranite in Batouri yielded an age of 620 Ma (U-Pb on zircon; Asaah VA, 2014). As per the deformation

chronology of Li XH et al. (2017) and Ngako V et al. (2008), this corresponds to a D_1 - D_2 event interpreted as the peak of the granulitic metamorphism in the south domain of the CAFB. In addition, transpressional shear zone-related gold mineralization is reported from the Tcholliré Shear Zone (TSZ), the Betaré Oya Shear Zone (BOSZ) and in subsidiary shear zones in the adjacent areas (Tchameni R et al., 2013; Asaah VA et al., 2014). In the TSZ, gold lode is associated

with NE-trending shear zones and subsidiary fractures related to dextral shear (Tchameni R et al., 2013). In Batouri and Dimako-Mboscosso adjacent to the BOSZ, the lode gold deposit is found in quartz veins in steeply dipping brittle-ductile shear zones in foliated biotite meta-granite, wall rock, and in NNW to WNW trending faults inside the alkali-feldspar granites (Vishiti A et al., 2017). However, the gold-bearing quartz veins are more frequently recorded within the




Fig. 7. Field photographs of faults associated to $N40^{\circ}E$ to $N50^{\circ}E$ shear zones. Location of snaps for the below sub-figures are indicated in Fig. 5. a—a network of fractures ($N70^{\circ}E$ and $N45^{\circ}E$) filled up with quartz cross cutting the $N45^{\circ}E$ striking mylonitic band in the area of Fotsi Sud. Notice that the $N70^{\circ}E$ striking fracture cross-cut that of $N45^{\circ}E$. b—sub-vertical E-W striking fault plane filled up with sheet-like granite and basalt in the northern Fotsi. c—W-E striking sub-vertical sheet-like granite transposing $N30^{\circ}E$ striking mylonitic foliation in the northern Fotsi. d—NW-SE trending sheet-like granite in the Banwa town. Fractures can be found after zooming the image. e—parallel NNE-trending sheet-like granite cross cutting the $N30^{\circ}E$ trending mylonitic gneiss. On zooming the image one sees few P-planes defining the slip sense. The Y-plane is parallel to the half arrows. f— $N45^{\circ}E$ striking fracture associated with $N45^{\circ}E$ striking mylonitic gneiss in the Kékem area.




Fig. 8. Stress regime and shear planes in the study area. This is also as per Passchier CW and Trouw RAJ (2005).

NE-trending shear zone, which is about 13 km long and 100 m to 1 km wide (Vishiti A et al., 2017). Minor anastomosing shears with gold-bearing quartz veins trend NNE-SSW, NE-SW, and ENE-WSW with steep dip towards NW (Asaah VA et al., 2014).

The geometric distribution of these gold-bearing quartz veins is consistent with the Riedel shear model following a dextral slip along the main NE-ENE striking Betare Oya Shear Zone (Vishiti A et al., 2017). This gold bearing quartz veins system, which cross cuts variably deformed granitoids, formed during the Pan-African orogeny in a volcanic-arc setting, are composed principally of quartz \pm calcite and Mg-rich ankerite. This mineral assemblage is associated with major hydrothermal ore minerals such as arsenopyrite, chalcopyrite, pyrrhotite and galena.

The gold mineralization associated with transpression has been addressed widely (e.g., Araújo FJOD and Kuyumjian, 2000; Araújo FJOD et al., 2005; Tchameni R et al., 2013; Groves DI et al., 2018; Silva FW et al., 2018). Both the BOSZ and TSZ, parallel to CCSZ, are simple shear-dominated zones of transpression (Njonfang E et al., 2008; Kankeu B et al., 2009; Bella Nké BE et al., 2018) of lithospheric-scale (Njonfang E et al., 2008) constrained at 570–552 Ma (Tchaptchet TD et al., 2009). Those are also the zones of fluid flow (Groves DI et al., 1988) sourced from (1) asthenospheric upwelling (Bierlein PF et al., 2006), (2) dehydration metamorphic reaction of carbonaceous metasedimentary rocks, and/or (3) fluid released during felsic and intermediate magma crystallization (Silva FW et al., 2018). In these shear zone, as the deformation intensifies, the transverse shortening component increases the strike-slip that deviates from the simple shear. The strike slip deformation originated vertical mylonitic foliations with sub-horizontal lineations on them. Plastic deformation and dynamic recrystallization of sub-grain modified the permeability of the deforming rocks resulting in fluid accumulation leading to micro-cracks (Zhu J et al., 2014) and favorable environment for gold deposits (Zhu J et al., 2014).

When the transpression/shear zone is associated with genetically linked subsidiary shear zones, the latter are referred also as the first-order/second-order brittle planes. The

first order shear zone is characterized by higher temperature commonly containing gold dissolved as reduced sulphide with respect to the gold occurring at the secondary fractures. Contrasting fluid-pressure and/or temperature promotes fluid flow and gold deposition. This hypothesis can be applied for the gold deposits in Cameroon because the last dextral shear of the transpressional tectonics occurring at high temperature (Njonfang E et al., 2008) is susceptible to rehomogenize gold in 620 Ma-aged syn-collisional granitoids into subsidiary Riedel shears along the main shear zone. This hypothesis matches with Araújo FJOD and Kuyumjian RM (2000). As per Araújo FJOD and Kuyumjian RM (2000), the gold deposits in the NE Brazil are probably due to the reactivation during D₃ strike-slip deformation, which remobilized gold of the Pre S-granitoids into Riedel's fractures kinematic models.

Summarily, the main gold mineralization is presumably associated with the last D₃ transpressional shear about 570–552 Ma. This is even though the gold-bearing 640–620 Ma syn-collisional granitoids is recorded in Batouri (summary from Asaah VA et al., 2014). These transpression shear zone-related gold mineralizations are probably coeval with the reactivation of the early structures during the last collisional activities amongst the Congo Craton, West African Congo Craton and Saharan Metacraton, as suggested by Ngako V et al. (2008).

7. Potential targets for gold exploration within the CAFB

As discussed, crustal environment favorable for gold deposits is apparently controlled by the structural setting and the lithology. Lithospheric-scale transpression/shear zones characterized by the association of pure- and simple shear followed by mylonization can result in fluid focusing and gold deposition. Greenschist facies metasedimentary rocks and high K alkaline to alkaline granite are gold hosting rocks, suggesting gold-enriched fluid (Asaah VA et al., 2014) to be present in the past. These geological features are reported from Kékem by Tchaptchet TD et al. (2009) and Tcheumenak Kouémo J et al. (2014). According to these authors, the Kékem area forms the SW prolongation of the CCSZ. In addition, metasedimentary rocks experienced a clockwise P-T path with a retrograde loop characterized by an early D₁, D₂ deformations during the upper amphibolite facies metamorphism and mylonitization between 570–552 Ma. These metamorphic rocks are intruded by Pan-African felsic to intermediate magmatic rocks (Tchaptchet TD et al., 2009). Both events can generate gold-enriched fluid (Tomkins GA et al., 2013; Silva FW et al., 2018) that flow through the fractures. Also, the reactivation process ascribed to D₃ event (Tcheumenak Kouémo J et al., 2014), which operates at high temperature is able to rehomogenize the gold-enriched fluid that move to low strain zones i.e., fracture with Riedel kinematic model (this study). This argument is strengthened by numerous quartz veins associated to fractures that connote fluid activity (Groves DI et al., 1988). These quartz veins hold gold grains / lode as in Betaré-Oya.

The similarities in lithological association, metamorphism facies (greenschist facies) and structural evolution between the Batouri and Betaré Oya area where gold-bearing quartz veins are reported and the Kékem area that forms the SW of the CCSZ, suggests that the CCSZ as well as the subsidiary shear fractures in the adjacent areas, are crustal environment favorable for gold deposits. Accordingly, the central domain of the CAFB bounded by transpression/shear zone associated with subsidiary Riedel's shear fractures kinematic model can be the potential target for further gold exploration.

8. Conclusions

(i) The Kékem area forms the SW prolongation of the CCSZ. Evolution of this segment of the CCSZ is characterized by an early D_1 crustal thickening, followed by D_2 sinistral shear and a D_3 dextral simple shear-dominated transpression and then a D_4 brittle deformation. This brittle deformation, presumably no older than 550 Ma, is the onset of the extensional tectonics as illustrated by associated granites and tholeiitic basalts.

(ii) The main gold mineralization is associated with 570–552 Ma D_3 simple shear-dominated transpression (Tcholliré and Betaré Oya shear zone) and secondary fractures of their adjacent regions. However, syn-collisional (syn- D_2 granitoids) gold lodes exist in Batouri. The reactivation of the early structures during late collision activities through simple shear rehomogenized the syn-collisional granitoid-related gold.

(iii) The deep-sourced fluid might have mixed with the component released from dehydration metamorphic reaction and felsic magma crystallization. These CO_2 -enriched fluids contain gold dissolved as reduced sulphide at high temperature, and deposited gold in dilation zone where mylonitization operated under greenschist to amphibolite facies conditions.

(iv) The similarity in structural evolution and lithologies of the gold-bearing Betaré Oya transpressional shear zone with that the Kékem area that forms the SW prolongation of the CCSZ, and suggests a regional setting controlling the gold mineralization within the CAFB. Therefore, the CCSZ as well as its adjacent areas with secondary fractures could be a suitable spot for gold exploration.

CRedit authorship contribution statement

Tchaptchet Tchato is the one who has carried out field study, conceived and initiated the paper in collaboration with Professor Soumyajit Mukherjee. In addition, the latter reorganise the paper, redraw some figure with corel draw software according to editor's requirements. Numbem has actively participated in writing the manuscript and has highly improved the english language quality. Ngamy has contributed in the methodology section and has produced original figures for the paper. Tchouankoue upgrade the manuscript by reviewing it.

Declaration of competing interest

There are no conflicts of interest to declare.

Acknowledgement

The Chief Editor, the Managing Editor, the Handling Editor (Xi-jie Chen), and the two anonymous reviewers are thanked. CPDA Grant (IIT Bombay) supported SM.

References

- Ahlgren SG. 2001. The nucleation and evolution in Riedel shear zones as deformation bands in porous sandstone. *Journal of Structural Geology*, 23(8), 1203–1214. doi: [10.1016/S0191-8141\(00\)00183-8](https://doi.org/10.1016/S0191-8141(00)00183-8).
- Akam JM, Mvondo OJ, Assatse TW, Owona S, Olinga JB, Messi OEJ, Ntomba S. 2014. Apport des images landsat 7 ETM+ à l'étude structurale du socle archéen de Sangmélina (Sud-Cameroun). *Revue Française de Photogrammétrie et de Télédétection*, 206, 15–25 (in French with English abstract). doi: [10.52638/rfpt.2014.119](https://doi.org/10.52638/rfpt.2014.119).
- Arau'jo CNM, Vasconcelos MP, da Silva FCA, de Sá EFJ, Sá JM. 2005. $^{40}Ar/^{39}Ar$ geochronology of gold mineralization in Brasileiro strike-slip shear zones in the Borborema province, NE Brazil. *Journal of South American Earth Sciences*, 19(4), 445–460. doi: [10.1016/j.jsames.2005.06.009](https://doi.org/10.1016/j.jsames.2005.06.009).
- Araújo FJOD, Kuyumjian RM. 2000. Structurally-controlled gold occurrences in the southern Brasília fold-thrust belt of central and SE-goias. *Revista Brasileira de geociencias*, 30(2), 289–292. doi: [10.25249/0375-7536.2000302289292](https://doi.org/10.25249/0375-7536.2000302289292).
- Archanjo CJ, Viegas LGF, Hollanda MHBM, Souya CL, Liu D. 2013. Timing of the HT/LP transpression in the Neoproterozoicserido Belt (Borborema Province, Brazil): Constraints from U-Pb (SHRIMP) geochronology and implications for the connections between NE Brazil and West Africa. *Gondwana Research*, 23(2), 701–714. doi: [10.1016/j.gr.2012.05.005](https://doi.org/10.1016/j.gr.2012.05.005).
- Aasaah VA, Zoheir B, Lehmann B, Frei D, Burgess R, Suh EC. 2014. Geochemistry and geochronology of the – 620 Ma gold-associated Batouri granitoids, Cameroon. *International Geology Review*, 57(11–12), 1485–1509.
- Bella Nké BE, Njanko T, Mamtani MA, Njonfang E, Rochette P. 2018. Kinematic evolution of the Mbakop Pan-African granitoids (western Cameroon –Domain): An integrated AMS and EBSD approach. *Journal of Structural Geology*, 111, 42–63. doi: [10.1016/j.jsg.2018.03.006](https://doi.org/10.1016/j.jsg.2018.03.006).
- Bierlein PF, Groves ID, Goldfarb JR, Dubé B. 2006. Lithospheric controls on the formation of provinces hosting giant orogenic gold deposits. *Mineralium Deposita*, 40, 874–886. doi: [10.1007/s00126-005-0046-2](https://doi.org/10.1007/s00126-005-0046-2).
- Bouyo MH, Penaye J, Barbey P, Toteu SF, Wandji P. 2013. Petrology of high pressure granulite facies metapelites and metabasites from Tcholliré and Banyo regions: Geodynamic implication for the central African fold belt (CAFB) of north-central Cameroon. *Precambrian Research*, 224, 412–413. doi: [10.1016/j.precamres.2012.09.025](https://doi.org/10.1016/j.precamres.2012.09.025).
- Bouyo MH, Zhao Y, Penaye J, Zhang SH, Njel UO. 2015. Neoproterozoic subduction-related metavolcanic and metasedimentary rocks from the Rey Bouba Greenstone Belt of north-central Cameroon in the Central African Fold Belt: New insights into a continental arc geodynamic setting. *Precambrian Research*, 261, 40–53. doi: [10.1016/j.precamres.2015.01.012](https://doi.org/10.1016/j.precamres.2015.01.012).
- Caby R. 1989. Precambrian terranes of Benin-Nigeria and Northeast Brazil and the late Paleogeographical reconstruction of the Pan-African / Brasileiro orogen: Closure of an oceanic domain of intracontinental convergence between major blocks. *Precambrian*

- Research, 69, 327–344.
- Dasgupta S, Mukherjee S. 2019. Remote sensing in lineament identification: Examples from western India, In: Billi A, Fagereng A. (eds.) Problems and Solutions in Structural Geology and Tectonics. Developments in Structural Geology and Tectonics book series, Series Editor: Mukherjee S, Elsevier.
- Davis GH, Bnmp AP, Garcia PE, Ahlgren SG. 2000. Conjugate Riedel deformation band shear zones. *Journal of Structural Geology*, 22(2), 169–190. doi: [10.1016/S0191-8141\(99\)00140-6](https://doi.org/10.1016/S0191-8141(99)00140-6).
- Dumort JC. 1968. Notice explicative de la carte géologique de reconnaissance du Cameroun au 1/500000. Feuille Douala Ouest. Direction des Mines et Géologie du Cameroun. 69 pp. (in French)
- Ganwa AA, Frisch W, Mvondo OJ, Njom B. 2007. Relationships between the parameters of geomorphology and structural features in the Pan-African fold belt of Cameroon. Example of Kombé II-Mayabo Area. *Journal of Engineering and Applied Sciences*, 2(2), 336–341.
- Ganwa AA, Klotzli UST, Hauzenberger C. 2016. Evidence for Archean inheritance in the pre-Pan-African crust of Central Cameroon: Insight from zircon internal structure and LA-MCICP-MS U-Pb ages. *Journal of African Earth Sciences*, 120, 12–22. doi: [10.1016/j.jafrearsci.2016.04.013](https://doi.org/10.1016/j.jafrearsci.2016.04.013).
- Groves DI, Goldfarb JR, Gebre-Mariam M, Hagemann SG, Robert F. 1988. Orogenic gold deposits: A proposed classification in the context of their crustal distribution and relationship to other gold deposit types. *Ore Geology Reviews*, 13(1–5), 7–27.
- Groves DI, Santosh M, Goldfarb JR, Zhang L. 2018. Structural geometry of orogenic gold deposits: Implications for exploration of world class and giant deposits. *Geoscience Frontiers*, 9(4), 1163–1177. doi: [10.1016/j.gsf.2018.01.006](https://doi.org/10.1016/j.gsf.2018.01.006).
- Kankeu B, Greiling RO, Nzenti JP. 2009. Pan-African strike-slip tectonics in eastern Cameroon-Magnetic fabrics (AMS) and structure in the Lom basin and its gneissic basement. *Precambrian Research*, 174(3–4), 258–272. doi: [10.1016/j.precamres.2009.08.001](https://doi.org/10.1016/j.precamres.2009.08.001).
- Kwekam M, Affaton P, Bruguier O, Liégeois JP, Harmann G, Njonfang E. 2013. The Pan-african Kékem gabbro-norite (West-cameroon), U-Pb zircon age, geochemistry and Sr-Nd isotopes: Geodynamic implication for the evolution of the central African fold belt. *Journal of African Earth Sciences*, 84, 70–88. doi: [10.1016/j.jafrearsci.2013.03.010](https://doi.org/10.1016/j.jafrearsci.2013.03.010).
- Li XH, Chen Y, Tchouankoué JP, Liu CZ, Li J, Ling XX, Tang GQ, Liu Y. 2017. Improving geochronological framework of the Pan-African orogeny in Cameroon: New SIMS zircon and monazite U-Pb age constraints. *Precambrian Research*, 294, 307–321. doi: [10.1016/j.precamres.2017.04.006](https://doi.org/10.1016/j.precamres.2017.04.006).
- Milési JP, Toteu SF, Deschamps Y, Feybesse JL, Lerouge C, Cocherie A, Penaye J, Tchameni R, Moloto A-Kenguemba G, Kampunzu, HAB, Nicol N, Duguey E, Leistel JM, Saint-Martin M, Ralay F, Henry C, Bouchot V, Doumnang Mbaigane JC., Kanda Kula V, Chene F, Montheil J, Boutin P, and Cailteux J. 2006. An overview of the geology and major ore deposits of Central Africa: Explanatory note for the 1:4000000 map 'Geology and major ore deposits of Central Africa'. *Journal of African Earth Sciences*, 44, 571–595. doi: [10.1016/j.jafrearsci.2005.10.016](https://doi.org/10.1016/j.jafrearsci.2005.10.016).
- Misra AA, Bhattacharya G, Mukherjee S, Bose N. 2014. Near N-S paleo-extension in the western Deccan region in India: Does it link strike-slip tectonics with India-Seychelles rifting? *International Journal of Earth Sciences*, 103, 1645–1680. doi: [10.1007/s00531-014-1021-x](https://doi.org/10.1007/s00531-014-1021-x).
- Misra S, Mandal N, Chakraborty C. 2009. Formation of Riedel shear fractures in granular materials: Findings from analogue shear experiments and theoretical analyses. *Tectonophysics*, 471(3–4), 253–259.
- Mukherjee S. 2013. Channel flow extrusion model to constrain dynamic viscosity and Prandtl number of the Higher Himalayan Shear Zone. *International Journal of Earth Sciences*, 102, 1811–1835. doi: [10.1007/s00531-012-0806-z](https://doi.org/10.1007/s00531-012-0806-z).
- Mukherjee S, Khonsari MM. 2018. Inter-book normal fault-related shear heating in brittle bookshelf faults. *Marine and Petroleum Geology*, 97, 45–48. doi: [10.1016/j.marpetgeo.2018.06.029](https://doi.org/10.1016/j.marpetgeo.2018.06.029).
- Mvondo H, den Brok SWJ, Ondo JM. 2003. Evidence of symmetric extension and exhumation of the Yaoundé nappe (Pan-African fold belt, Cameroon). *Journal of African Earth Sciences*, 36(3), 215–231. doi: [10.1016/S0899-5362\(03\)00017-4](https://doi.org/10.1016/S0899-5362(03)00017-4).
- Mvondo H, Owona S, Ondo JM, Essono J. 2007. Tectonic evolution of the Yaoundé segment of the Neoproterozoic Central African Orogenic Belt in southern Cameroon. *Canadian Journal of Earth Sciences*, 44(4), 433–444. doi: [10.1139/e06-107](https://doi.org/10.1139/e06-107).
- Neawsuparp K, Charusiri P. 2004. Lineaments analysis determined from Landsat Data: Implication for tectonic features and mineral occurrences in Northern Loei Area, NE Thailand. *Science Asia*, 30, 269–278. doi: [10.2306/scienceasia1513-1874.2004.30.269](https://doi.org/10.2306/scienceasia1513-1874.2004.30.269).
- Ngako V, Njonfang E. 2011. Plates Amalgamation and Plate Destruction, the Western Gondwana History. In: Closson D (ed.). *Tectonics*. United Kingdom, INTECH, 358.
- Ngako V, Affaton P, Njonfang E. 2008. Pan-African tectonics in northern Cameroon: Implication for the history of western Gondwana. *Gondwana Research*, 14(3), 509–522. doi: [10.1016/j.gr.2008.02.002](https://doi.org/10.1016/j.gr.2008.02.002).
- Ngako V, Jégouzo P, Nzenti JP. 1991. Le Cisaillement Centre Camerounais. Rôle structural et géodynamique dans l'orogénèse panafricaine. *Comptes Rendus de l'Académie des Sciences*, 313, 457–463 (in French with English abstract).
- Ngamy KA, Owona S, Tchakounté JN, Mvondo Ondo J. 2016. Fabriques en aplatissement et constriction des granitoïdes de Nga Mbappé et de Yoro au nord du Groupe de Yaoundé (CPNE, Cameroun): témoins de la mise en place de granites syn-tectoniques au cours de l'orogénèse Pan-Africaine. *Journal of Comunicações Geológicas*, 103, 1, 5–16.
- Njome MS, Suh CE. 2005. Tectonic evolution of the Tombel graben basement, southwestern Cameroon. *Episodes*, 28(1), 37–41. doi: [10.18814/epiiugs/2005/v28i1/004](https://doi.org/10.18814/epiiugs/2005/v28i1/004).
- Njonfang E, Ngako V, Moreau C, Affaton P, Diot E. 2008. Restraining bends in high temperature shear zones: The Central Cameroon Shear Zone, central Africa. *Journal of African Earth Sciences*, 52(1–2), 9–20. doi: [10.1016/j.jafrearsci.2008.03.002](https://doi.org/10.1016/j.jafrearsci.2008.03.002).
- Nouthogwé TC, Koumetio F, Manguelle-Dicoum E. 2010. Structural features of South –Adamawa (Cameroon) inferred from magnetic anomalies: Hydrogeological implications. *Compte Rendus Geoscience*, 342(6), 467–474. doi: [10.1016/j.crte.2010.03.004](https://doi.org/10.1016/j.crte.2010.03.004).
- Owona S, Schulz B, Ratschbacher L, Mvondo OJ, Ekodeck G-E, Tchoua MF, Affaton P. 2010. Pan-African metamorphic event in the southern Yaoundé Group (Oubanguide Complex, Cameroon) as revealed by EMP-monazite and thermobarometry of garnet metapelites. *Journal of African Earth Sciences*, 59, 125–139.
- Passchier CW, Trouw RAJ. 2005. *Microtectonics*, 2nd edition. Berlin, Springer-Verlag, Heidelberg.
- Penaye J, Toteu SF, Tchameni R, Van Schmus WR, Tchakounté J, Ganwa A, Miyem D, Nsifa EN. 2004. The 2.1 Ga west Central African Belt in Cameroon: Extension and evolution. *Journal of African Earth Sciences*, 39(3–5), 159–64. doi: [10.1016/j.jafrearsci.2004.07.053](https://doi.org/10.1016/j.jafrearsci.2004.07.053).
- Penaye J, Toteu SF, Van Schmus WR, Nzenti JP. 1993. U-Pb and Sm-Nd preliminary geochronologic data on the Yaoundé series, Cameroon: Re-interpretation of the granulitic rocks as the suture of a collision in the Central african belt. *Comptes Rendus de l'Académie des Sciences Paris*, 317, 789–94.
- Schreurs G, Colleta B. 1998. Analogue modelling of faulting in zones of continental transpression and transtension. In: Holdsworth RE,

- Strachn RA, Dewey JF (eds) Continental Transpressional and Transtensional Tectonics. Geological Society, London, Special Publications, 135, 59–79.
- Silva FW, Rosière AC, Buhn B. 2018. The shear zone-related gold mineralization at the Turmalina deposit, Quadrilatero Ferrifero, Brazil: Structural evolution and the two stages of mineralization. *Mineralium Deposita*, 54, 347–368. doi: <https://doi.org/10.1007/s00126-018-0811-7>.
- Simeni NAW, Tchaptchet TD, Ngo Belnoun RN, Tchouankoué JP, Ganwa AA. 2017. Structural relationship between brittle deformation and Paleozoic to Mesozoic basalt dykes in the Precambrian basement of the southern continental part of the Cameroon Volcanic Line. *International Journal of Geoscience*, 8(3), 318–331. doi: [10.4236/ijg.2017.83016](https://doi.org/10.4236/ijg.2017.83016).
- Som Mbang MC, Basseka AC, Kamguia J, Etamè J, Tchoukeu NDC, Mouzong PM. 2018. Mapping of deep tectonic structures of central and southern Cameroon by an interpretation of surface and satellite magnetic data. *International Journal of Geophysics*, 5845670. doi: <https://doi.org/10.1155/2018/5845670>.
- Stendal H, Toteu SF, Frei R, Penaye J, Njel UO, Bassahak J, Nni J, Kankeu B, Ngako V, Hell JV. 2006. Derivation of detrital rutile in the Yaounde region from the Neoproterozoic Pan-African belt in southern Cameroon (Central Africa). *Journal of African Earth Sciences*, 44(4–5), 443–458. doi: [10.1016/j.jafrearsci.2005.11.012](https://doi.org/10.1016/j.jafrearsci.2005.11.012).
- Suh CE, Lehmann B, Mafany TG. 2006. Geology and geochemical aspects of the lode gold mineralization at Dimako-Mboscorro, SE Cameroon. *Geochemistry: Exploration, Environment, Analysis*, 6(4), 295–309. doi: [10.1144/1467-7873/06-110](https://doi.org/10.1144/1467-7873/06-110).
- Takodjou WJD, Ganno S, Afahwie NA, Nomo NE, Mvondo OJ, Nzenti JP. 2016. Use of Landsat 7 ETM+Data for the geological structure interpretation: Case study of the Ngoura-Colomines Area, Eastern Cameroon. *Journal of Geosciences and Geomatics*, 4(3), 61–72.
- Tchameni R, Doumnang JC, Deudibaye M, Branquet Y. 2013. On the occurrence of gold mineralization in the Pala Neoproterozoic formations, South-Western Chad. *Journal of African Earth Sciences*, 84, 36–46. doi: [10.1016/j.jafrearsci.2013.03.002](https://doi.org/10.1016/j.jafrearsci.2013.03.002).
- Tchaptchet TD, Schulz B, Nzenti JP. 2009. Electron microprobe monazite dating and thermobarometry of the Neoproterozoic metamorphic events in the Kékem area, central Africa fold of Cameroon. *Neues Jahrbuch für Mineralogie Abhandlungen*, 186(1), 95–109.
- Tchaptchet TD. 2011. Geology of the Kékem area (Cameroon central domain): Metaamorphic petrology, P-T-t path, EMP (monazite), LA-ICPMS (zircon) dating and implications for the geodynamic evolution of the Pan-African North Equatorial fold belt. Université de Yaoundé I, Thèse doctorat PhD, Yaoundé 109 pages.
- Tchaptchet TD, Wambo NAS, Kouamo NAK, Tchouankoué JP, Cucciniello C. 2017. Geology, mineralogy and geochemistry of the Kékem dyke swarm (Western Cameroon): Insights into Paleozoic–Mesozoic magmatism and geodynamic implications. *Comptes Rendus Geoscience*, 349(4), 175–85. doi: [10.1016/j.crte](https://doi.org/10.1016/j.crte.2017.02.005).
- Tcheumenak Kouémo J, Njanko T, Kwékam M, Naba S, Nké BEB, Sandjo AFY, Fozing EM, Njonfang E. 2014. Kinematic evolution of the Fodjomekwet-Fotouni Shear Zone (West-Cameroon): Implications for emplacement of the Fomopéa and Bandja plutons. *Journal of African Earth Sciences*, 99, 261–275. doi: [10.1016/j.jafrearsci.2014.07.018](https://doi.org/10.1016/j.jafrearsci.2014.07.018).
- Tchuimegnie Ngongang NB, Kamgang P, Chazot G, Agranier A, Bellon H, Nonnotte P. 2015. Age, geochemical characteristics and petrogenesis of Cenozoic intraplate alkaline volcanic rocks in the Bafang region, West Cameroon. *Journal of African Earth Sciences*, 102, 218–232. doi: [10.1016/j.jafrearsci.2014.10.011](https://doi.org/10.1016/j.jafrearsci.2014.10.011).
- Tchakounté J, Eglinger A, Toteu SF, Zeh A, Nkoumbou C, Mvondo-Ondoa J, Penaye J, de Wit M, Barbey P. 2017. The Adamawa-Yadé domain, a piece of Archaean crust in the Neoproterozoic Central African Orogenic Belt (Bafia area, Cameroon). *Precambrian Research*, 299, 210–229.
- Tchakounté J. 2020. Tectonique, Géochronologie et Pétrogenèse du massif de Bapé et de son encaissant (Bafia, Centre-Cameroun): implications géodynamiques pour l'évolution crustale de la bordure sud du bloc Adamaoua-Yadé. Université de Yaoundé I, Thèse doctorat/PhD, Yaoundé, 215 pages.
- Tomkins GA. 2013. On the source of orogenic gold. *Geology*, 41(12), 1255–1256. doi: [10.1130/focus122013.1](https://doi.org/10.1130/focus122013.1).
- Toteu SF, Penaye J, Djomani YP. 2004. Geodynamic evolution of the Pan-African belt in Central Africa with special reference to Cameroon. *Canadian Journal of Earth Sciences*, 41(1), 73–85. doi: [10.1139/e03-079](https://doi.org/10.1139/e03-079).
- Toteu SF, van Schumus WR, Penaye J, Michard A. 2001. New U-Pb and Sm-Nd data from North-Central Cameroon and its bearing on the pre-pan African history of Central Africa. *Precambrian Research*, 108(1–2), 45–73.
- Toteu SF, Penaye J, Deloule E, van Schmus WR, Tchameni R. 2006. Diachronous evolution of Volcano–sedimentary basins north of the Congo Craton: Insights from U-Pb ion microprobe dating of zircons from the Poli, Lom and Yaoundé Groups (Cameroon). *Journal of African Earth Sciences*, 44(4–5), 428–442.
- Vishiti A, Suh CE, Lehmann B, Shemang EM, Lionel N, Ngome J, Nshanji NJ, Chinjo FE, Mongwe OY, Petersen S. 2017. Mineral chemistry, bulk rock geochemistry, and S - isotope signature of lode - gold mineralization in the Bétaré Oya gold district, south - east Cameroon. *Geological Journal*, 53(6), 2579–2596.
- Vishiti A, Etame J and Suh CE. 2019. Features of gold bearing quartz veins in an artisanal mining–dominated terrain, Batouri gold district, Eastern region of Cameroon. *Episodes*, 42(3), 199–212. doi: [10.18814/epiiugs/2019/019016](https://doi.org/10.18814/epiiugs/2019/019016).
- Zhu J, Li Z, Lin G, Zeng Q, Zhou Y, Yi J, Gong G, Chen G. 2014. Numerical simulation of mylonitization and structural controls on fluid flow and mineralization of the Hetai gold deposit, west Guangdong, China. *Geofluids*, 14, 221–233. doi: [10.1111/gfl.12069](https://doi.org/10.1111/gfl.12069).